



## Original Research Article

# Neurophysiology of pain education knowledge, pain disability, patient satisfaction and central sensitization in chronic musculoskeletal pain

Thangamani Ramalingam<sup>1,\*</sup>, Pooja Desai<sup>1</sup>, Dhruvi Ghoghari<sup>1</sup>, Vidhi Jethva<sup>1</sup>, Rushvi Shah<sup>1</sup>

<sup>1</sup>Dept. of Physiotherapy, Sarvajani College of Physiotherapy, Surat, Gujarat, India



## ARTICLE INFO

## Article history:

Received 14-12-2022

Accepted 29-12-2022

Available online 23-01-2023

## Keywords:

Chronic musculoskeletal pain  
Neurophysiology of pain education knowledge  
Central sensitization  
Pain disability  
Patient satisfaction

## ABSTRACT

**Introduction:** Chronic musculoskeletal pains are multifaceted, and Central sensitization is a potential pathophysiological mechanism underlying a group of chronic musculoskeletal pain disorders. Neurophysiology of pain education knowledge and patient satisfaction levels of chronic musculoskeletal pain subjects may contribute to central nervous system sensitization. Hence, the aim of the study was to evaluate the impact of neurophysiology of pain education knowledge and patient satisfaction levels on central sensitization in large population of patient with chronic musculoskeletal pain.

**Methods Subjects:** The study included 200 chronic musculoskeletal pain subjects that persisted more than 3 months with average age of  $43.93 \pm 13.62$ .

**Study Design and Sampling:** A cross-sectional study used non probability sampling.

**Outcome Measures:** Neurophysiology of pain Questionnaire (NPQ) to know the conceptualization of pain, mood rating scale (MRS) to measure patient's mood fluctuation because of pain, pain disability scale (PDS) for evaluating patients ability to perform certain activity, central sensitization inventory (CSI) to measure nervous system sensitization and patients satisfaction scale (PSS) to understand patient's satisfaction towards the treatment and health care provider were used.

**Statistical Analysis:** Descriptive and correlation analyses were used for analysis.

**Results:** The correlation analyses showed that patient disability scale negatively correlated with mood scale and positively correlated with the age, duration of the condition, impact of pain on ADL and central sensitization. And there was a positive correlation between patient satisfaction and impact of pain on ADL.

**Conclusion:** The age, mood, duration of the condition and pain disability were the factors behind central sensitization in patients with chronic musculoskeletal pain. And the neuro physiology of pain knowledge had impact on pain disability and patient satisfaction.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Pain, an unpleasant sensory and emotional experience is one of the most burdensome problems of recent time which demands intervention and challenges the health care system.<sup>1,2</sup> In clinical practice, many consider persistent or recurrent pain for  $\geq 3$  months to be a useful definition of the

musculoskeletal pain.<sup>3</sup> The new international classification of diseases (ICD-11) categorization identifies chronic musculoskeletal pain in two forms (CMP), the primary and secondary. CMP conditions are multifaceted, and approximately 20% of the adult population lives with severe chronic pain with higher prevalence in women and in lower income groups. Chronic pain is influenced by and interacts with physical, emotional, psychological, and social factors, and a bio psychosocial framework is increasingly

\* Corresponding author.

E-mail address: [atramalingam@gmail.com](mailto:atramalingam@gmail.com) (T. Ramalingam).

applied in clinical practice<sup>4,5</sup> chronic musculoskeletal pain is the main contributor to increased suffering in daily activities, drug consumption and disability leading to more health cost worldwide.<sup>6-8</sup> Central sensitization is a neurophysiological phenomenon that is adaptive, activity dependent, and dynamic and has recently been recognized as a potential pathophysiological mechanism underlying a group of chronic pain disorders. In addition, the term 'central' may refer to ipsilateral sensitization, segmental sensitization, extraterritorial and generalized widespread sensitization.<sup>9-11</sup> Central sensitization encompasses various related dysfunctions within the central nervous system, all contributing to increased responsiveness to a variety of stimuli including mechanical pressure, chemical substances, light, sound, cold, heat, stress, and electrical stimulation.<sup>12</sup> Pain represents an experience that is influenced not only by sensation, but also by context and prior experience. The experience of pains distinct from nociception describes afferent neural activity transmitting sensory information about stimuli that have the potential to cause tissue damage.<sup>13</sup> Patient satisfaction and preference to be considered important while making the treatment decision along with patient information and involvement in the process.<sup>14</sup> Pain neuroscience education(PNE), is a cognitive intervention intended to modify the inappropriate beliefs and maladaptive behaviours of chronic pain patients to decrease pain and disability.<sup>15</sup> PNE is believed to desensitize the central nervous system and Combining that to therapeutic exercise resulted in significantly better results for participants with chronic pain, with a large effectsize, compared to therapeutic exercise alone.<sup>16</sup> Hence, the aim of the study was to find out the correlation of neurophysiology of pain education level, pain disability, central sensitization and patient satisfaction in patient with chronic musculoskeletal pain.

## 2. Materials and Methods

### 2.1. Participants

A cross-sectional study was done using non probability sampling of 200 patients with chronic neck, back, shoulder, knee pain, pain more than 3 months and individuals who gave their consent only included.

### 2.2. Outcome measures

1. The Central Sensitization Inventory (CSI): The CSI consists of 25 questions and may be self-administered. Each question may be answered as follows: Never (0 points), rarely (1 point), Sometimes (2 points), Often (3 points), or Always (4 points). Total points reflect the severity of the central sensitization phenomenon.<sup>17</sup>
2. The Neurophysiology Pain Questionnaire(NPQ): Neurophysiology of pain education was devised to assess how an individual conceptualizes the biological

mechanisms that underpin his or her pain and scores ranging between 0(poor knowledge) to13(good knowledge).<sup>18</sup>

3. Pain Disability Questionnaire(PDQ): The items of the questionnaire are assessed on a 0–10 numeric rating scale in which 0 means no disability and 10 is maximum disability. The sum of the seven items equals the total score of the PDI, which ranges from 0 to 70, with higher scores reflecting higher interference of pain with daily activities.<sup>19</sup>
4. Short Assessment of Patient Satisfaction Scale (SAPS): It is a generic measure of care quality and patient satisfaction with 7 items on a five point Likert's scale, scores ranging from 0 (extremely dis satisfied) to 28 (extremely satisfied).<sup>20</sup>

### 2.3. Procedure

Consent was taken and the objectives of the study were explained to the subjects and instructions were given before filling the questionnaire and the data collection sheets.

## 3. Statistical Analysis

The categorical variables presented in percentages and numerical variables in terms of mean and standard deviations for the demographical and clinical outcomes. Descriptive and correlation analyses were used for analysis. Data were analyzed using IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp. Statistical significance was set at  $p < 0.05$  (Two tailed).

## 4. Results

The present study included 200 subjects with chronic musculoskeletal pain (with mean age  $43.94 \pm 13.63$ ) in the analysis in which 72 males and 128 females with pain duration of more than three months. The clinical characteristics data of all subjects along with descriptive statistics of the outcome measure are shown in the Table 1 &2. The correlation between the outcome measures and biopsychosocial factors are shown in the Table 3 &4.

## 5. Discussion

The present study included 200 patients with chronic musculoskeletal pain to find out the impact of neurophysiology of pain education, pain disability and patient satisfaction levels on central sensitization in chronic musculoskeletal pain. Patient satisfaction and Neurophysiology of pain education knowledge did not have any significant impact on central sensitization. But they are more significantly impacted by the perceived impact of activity of daily living in patients with chronic musculoskeletal pain disorders and further more age had low level of neurophysiology of pain knowledge in patients

**Table 1:** Bio psychosocial factors of participants (N=200)

Subject Characteristics	N	Percentage
Pain medication		
Yes	59	44.5
No	111	55.5
Onset of pain		
Sudden	39	19.5
Gradual	158	79.0
None	3	1.5
Mostly used posture		
Sitting	90	45
Standing	60	30
Walking	5	2.5
Sitting, Standing	40	20
All	5	2.5
Exercise per week		
1-2 days per week	51	25.5
3-4 days per wee	71	35.5
5 or more days	77	38.5
None	1	0.5
Coping mechanism		
Hot pack	40	20
Medicine	23	11.5
Rest	29	14.5
Cold pack	10	5
Other	98	49.5
Mood		
Very sad	15	7.5
Somewhat sad	77	38.5
Neutral	75	37.5
Somewhat happy	30	15
Very happy	3	1.5

**Table 2:** Descriptive statistics of outcome measures

Outcome measures	Mean± Standard Deviation
Duration of condition( in Months)	27.79±26.88
Work duration( in hours)	38.35±17.20
Impact of pain on ADL (0-10 point scale)	6.49±1.87
Neurophysiology of pain knowledge(NPQ)	4.44±1.79
Patient satisfaction(SAPS)	16.12±2.32
Central nervous system sensitization(CSI)	58.37±13.58
Pain disability(PDS)	69.37±25.70

with chronic musculoskeletal pain disorders.

The factors such as age, duration of pain, Mood and perceived impact of activity of daily living had a significant impact on central sensitization in patients with chronic musculoskeletal pain disorders which is similar to a study result from India. According to Lall D et al., central sensitization showed a significant association with pain discomfort, mood, and sleep <sup>21</sup> Pain disability had a significant association with central sensitization in patients with chronic musculoskeletal pain . According to Dennis C. Turk research study, the prevalence of depression as a comorbid psychological condition in chronic pain. <sup>21</sup> and

Tomoko Fujii study found high somatising tendency and higher pain ratings were associated with greater disability due to chronic musculoskeletal pain in different age groups. <sup>22,23</sup>

Neurophysiology of pain education knowledge did not have any significant impact on central sensitization. But significantly impacted by the perceived impact of activity of daily living and age in patients with chronic musculoskeletal pain. Recently in 2021, Kasper Bulow's research showed that the effects of Neurophysiology of pain education were moderate and statistically significant on pain intensity and psychological distress. <sup>24,25</sup> Perceived impact on activity of

**Table 3:** Correlation between the outcome measures


Outcome Measures	NPQ	SAPS	PDS	CSI
NPQ				
r:	1	-.175	-.215	.018
p:		.013	.002	.797
SAPS				
r:		1	0.129	-.110
p:			.068	.122
PDS				
r:			1	.418
p:				<.001
CSI				
r:				1
p:				

[r- Correlation; p-significance]

**Table 4:** Significant correlation between the outcome measures and bio psychosocial factors

Outcome Measures	Age	Impact of Pain on ADL	Duration of The Condition	Mood
r:	.215	.232	.211	-.291
CSI				
p:	.002	.001	.003	<.001
r:	-.181	-.212	—	—
NPQ				
p:	.010	.003	—	—
r:	—	.221	-	—
SAPS				
p:	—	.002	—	—
r:	.244	.560	—	-.262
PDS				
p:	.001	<.001	—	<.001

[r- Correlation; p-significance]



**Fig. 1:** Schematic hypothetical model conclusion of the study findings

daily living by the patients with chronic musculoskeletal pain had significant association with central sensitization, Neurophysiology of pain education knowledge, Pain disability and Patient satisfaction which is similar to the results of a study in older patients.<sup>26,27</sup>

The findings of the present study paved a way to the researchers to speculate a hypothetical model (Figure 1) to understand the relationships of Biopsychosocial factors, Neurophysiology of pain education knowledge, pain

disability, patient satisfaction and central sensitization in chronic musculoskeletal pain.

A limitation of this study is the relatively small size of the chronic musculoskeletal pain group and some sort of clinical trials could have helped to understand the study in more detailed manner with correlations between two sets of groups.

**6. Conclusion**

The age, mood, duration of the condition and pain disability levels were the factors behind central sensitization in patients with chronic musculoskeletal pain. And Levels of neuro physiological pain knowledge had impact on pain disability and patient satisfaction.

**7. Source of Funding**

None.


**8. Conflict of Interest**

The authors declare that there is no conflict of interest.

## References

- Loeser JD, Melzack R. Pain: an overview. *Lancet*. 1999;353(9164):1607–16.
- Linton SJ, Kienbacher T. Psychological Subgrouping to Assess the Risk for the Development or Maintenance of Chronic Musculoskeletal Pain: Is This the Way Forward? *Clin J Pain*. 2020;36(3):172–7.
- Mallick-Searle T, Sharma K, Toal P, Gutman A. Pain and Function in Chronic Musculoskeletal Pain-Treating the Whole Person. *J Multidiscip Healthc*. 2021;14:335–47. doi:10.2147/JMDH.S288401.
- Alagappan T. Physiotherapy practice in persistent chronic musculoskeletal pain. *Acad Lett*. 2021;2:1–5.
- Gerdle B, Ghafouri B, Ernberg M, Larsson B. Chronic musculoskeletal pain: review of mechanisms and biochemical biomarkers as assessed by the microdialysis technique. *J Pain Res*. 2014;7:313–9.
- El-Tallawy SN, Nalamasu R, Salem GI, Lequang JAK, Pergolizzi JV, Christo PJ. Management of Musculoskeletal Pain: An Update with Emphasis on Chronic Musculoskeletal Pain. *Pain Ther*. 2021;10(1):181–209.
- Wang C, Pu R, Ghose B, Tang S. Chronic Musculoskeletal Pain, Self-Reported Health and Quality of Life among Older Populations in South Africa and Uganda. *Int J Environ Res Public Health*. 2018;15(12):2806. doi:10.3390/ijerph15122806.
- Smith BE, Hendrick P, Smith TO, Bateman M, Moffatt F, Rathleff MS. Should exercises be painful in the management of chronic musculoskeletal pain? A systematic review and meta-analysis. *Brit J Sports Med*. 2013;51(23):1679–87.
- Van Griensven H, Schmid A, Trendafilova T, Low M. Central Sensitization in Musculoskeletal Pain: Lost in Translation? *J Orthop Sports Physical Ther*. 2020;50(11):592–6.
- Sanzarelli I, Merlini L, Rosa MA, Perrone M, Frugiuele J, Borghi R. Central sensitization in chronic low back pain: A narrative review. *J Back Musculoskelet Rehabil*. 2016;29(4):625–33.
- Arendt-Nielsen L, Morlion B, Perrot S, Dahan A, Dickenson A, Kress HG. Assessment and manifestation of central sensitisation across different chronic pain conditions. *Eur J Pain*. 2018;22(2):216–41.
- Nijs J, Leysen L, Vanlauwe J, Logghe T, Ickmans K, Polli A. Treatment of central sensitization in patients with chronic pain: time for change? *Expert Opin Pharmacother*. 2019;20(16):1961–70.
- Bourke JH, Langford RM, White PD. The common link between functional somatic syndromes may be central sensitisation. *J Psychosomatic Res*. 2015;78(3):228–64.
- Sippola S, Haijanen J, Viinikainen L, Grönroos J, Paajanen H, Rautio T. Quality of Life and Patient Satisfaction at 7-Year Follow-up of Antibiotic Therapy vs Appendectomy for Uncomplicated Acute Appendicitis: A Secondary Analysis of a Randomized Clinical Trial. *JAMA*. 2020;2020(4):283–9.
- King R, Robinson V, Elliott-Button HL, Watson JA, Ryan CG, Martin DJ. Pain Reconceptualisation after Pain Neurophysiology Education in Adults with Chronic Low Back Pain: A Qualitative Study. *Pain Res Manag*. 2018;p. 3745651. doi:10.1155/2018/3745651.
- Pardo GB, Girbés EL, Roussel NA, Izquierdo TG, Penick VJ, Martín DP. Pain Neurophysiology Education and Therapeutic Exercise for Patients With Chronic Low Back Pain: A Single-Blind Randomized Controlled Trial. *Arch Phys Med Rehabil*. 2018;99(2):338–47.
- Bid D, Neela S, Rathod P, Ramalingam T. Content Validity and Test-Retest Reliability of the Gujarati Version of the Central Sensitization Inventory. *J Integr Res Med*. 2016;7(5):18–24.
- Catley MJ, Connell NE, Moseley GL. How good is the neurophysiology of pain questionnaire? A Rasch analysis of psychometric properties. *J Pain*. 2013;14(8):818–45.
- Anagnostis C, Gatchel RJ, Mayer TG. The pain disability questionnaire: a new psychometrically sound measure for chronic musculoskeletal disorders. *Spine*. 2004;29(20):2290–302.
- Hawthorne G, Sansoni J, Hayes L, Marosszeky N, Sansoni E. Measuring patient satisfaction with health care treatment using the Short Assessment of Patient Satisfaction measure delivered superior and robust satisfaction estimates. *J Clin Epidemiol*. 2014;67(5):527–64.
- Turk DC, Fillingim RB, Ohrbach R, Patel KV. Assessment of Psychosocial and Functional Impact of Chronic Pain. *J Pain*. 2016;17(9):21–49.
- Fujii T, Oka H, Katsuhira J, Tonosu J, Kasahara S, Tanaka S. Disability due to knee pain and somatising tendency in Japanese adults. *BMC Musculoskeletal Dis*. 2018;19(1):23. doi:10.1186/s12891-018-1940-y.
- Louw A, Diener I, Butler DS, Puentedura EJ. The effect of neuroscience education on pain, disability, anxiety, and stress in chronic musculoskeletal pain. *Arch Phys Med Rehabil*. 2011;92(12):2041–56.
- Bülow K, Lindberg K, Vaegter HB, Juhl CB. Effectiveness of Pain Neurophysiology Education on Musculoskeletal Pain: A Systematic Review and Meta-Analysis. *Pain Med*. 2021;22(4):891–904.
- Lane E, Fritz JM, Greene T, Maddox D. The effectiveness of training physical therapists in pain neuroscience education on patient reported outcomes for patients with chronic spinal pain: a study protocol for a cluster randomized controlled trial. *BMC Musculoskelet Disord*. 2018;19(1):386. doi:10.1186/s12891-018-2269-2.
- Ritchie CS, Patel K, Boscardin J, Miaskowski C, Vranceanu AM, Whitlock E. Impact of persistent pain on function, cognition, and well-being of older adults. *J Am Geriatr Soc*. 2022;doi:10.1111/jgs.18125.
- Sugai K, Tsuji O, Matsumoto M, Nishiwaki Y, Nakamura M. Chronic musculoskeletal pain in Japan (the final report of the 3-year longitudinal study): Association with a future decline in activities of daily living. *J Orthop Surg*. 2017;25(3):2309499017727945. doi:10.1177/2309499017727945.

## Author biography

**Thangamani Ramalingam**, Faculty/Lecturer  <https://orcid.org/0000-0001-7429-1718>

**Pooja Desai**, Physiotherapy Graduates

**Dhruvi Ghoghari**, Physiotherapy Graduates

**Vidhi Jethva**, Physiotherapy Graduates

**Rushvi Shah**, Physiotherapy Graduates

**Cite this article:** Ramalingam T, Desai P, Ghoghari D, Jethva V, Shah R. Neurophysiology of pain education knowledge, pain disability, patient satisfaction and central sensitization in chronic musculoskeletal pain. *IP J Surg Allied Sci* 2022;4(4):137-141.