



## Original Research Article

# Clinical study of 104 cases of hollow viscus perforation with peritonitis in gastric and duodenal perforation at rural tertiary centre

Pranjal Tanter<sup>1,\*</sup>, Pratik Gond<sup>1</sup>, Shreya Pakkal<sup>1</sup>, Manideep Thippaarapu<sup>1</sup>, Kundankumar N Dandekar<sup>1</sup>

<sup>1</sup>Dept. of Surgery, Dr. Balasaheb Vikhe Patil Rural Medical College, Loni, Maharashtra, India



## ARTICLE INFO

## Article history:

Received 04-09-2023

Accepted 22-09-2023

Available online 13-10-2023

## Keywords:

Gastric perforation

Duodenal perforation

Graham's patch repair

Peritonitis

Hollow viscus perforation

Introduction

Peritonitis

Hollow viscus perforation

## ABSTRACT

**Background:** Acute abdomen with peptic ulcer perforation is a common cause of emergency admission and surgery. This is the study that documents the presentation and outcome of perforation management in DBVPRMC, Loni.

**Materials and Methods:** This is a prospective study of patients operated upon for perforation peritonitis between March 1, 2021, and March 23, 2023. A structured recorded after questionnaire containing patients demographics, operative findings, and outcome was discharge or death.

**Results:** There were 104 patients. 81 males and 23 females (M : F = 3.5 : 1). The age range was between 18 years and 95 years. The mean age was 48.99 years  $\pm$  SD 16.1 years. The ratio of gastric to duodenal perforation was 1.88 : 1. Perforation was the first sign of peptic ulcer disease in 62 (59.6%). Pneumoperitoneum was detectable with plain radiographs in 95 (91%) patients. 72 (69.2%) had Graham's Omentopexy. Death rate was 17.3%.

**Conclusion:** We note that gastric perforation is a more commoner disease in our environment. Perforation is often the first sign of peptic ulcer disease. We identify spicy food, alcoholism and NSAIDs intake amongst patient as a risk factor for perforation. Early presentation is associated with good prognosis.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Around 2-14% of instances of peptic ulcer disease result in peptic ulcer perforation, a potentially fatal complication.<sup>1,2</sup> This perforation causes the leakage of gastric contents into the peritoneal cavity and can be found either in the lesser curvature of the stomach or on the anterior surface of the duodenum.<sup>3</sup> One of the most frequent reasons for urgent hospitalisation and surgery for peptic ulcer disease is perforation.<sup>4,5</sup>

Princess Henrietta of England underwent the first clinical description of a perforated peptic ulcer in 1670.<sup>6</sup> Since then, this sickness has claimed the lives of several well-known

individuals.<sup>7</sup> The presentation may be dramatic, with quick start, acute pain that frequently radiates to the back, and fast developing symptoms.

In this traditional presentation, the patient may remember the precise moment of perforation, which was frequently in the early morning. Pain can occasionally be sneaky.

When the perforation is tiny and the contents slowly leak into the right iliac fossa through the right paracolic gutter, it can sometimes mimic an acute appendicitis<sup>8</sup> and present similarly in symptoms. The indications of perforation may be subtle or ambiguous in older patients or those who have impaired immune systems.<sup>9</sup>

The primary differential diagnosis is an abrupt exacerbation in a patient with known peptic ulcer disease, and the diagnosis is made with a high index of suspicion.<sup>10</sup>

\* Corresponding author.

E-mail address: [pranjaltanter@gmail.com](mailto:pranjaltanter@gmail.com) (P. Tanter).

Often, the diagnosis is established by the presence of air under the diaphragm on an erect chest radiograph. Up to 75%[12,33]<sup>11</sup> of erect chest radiographs contain this indicator. This indication, which can be shown on up to 75%[12,33]<sup>11</sup> of erect chest radiographs, depends on the size of the perforation and the time after the perforation was discovered. An erect lateral chest radiograph can increase pneumoperitoneum detection by 98% and can detect as 10cc of air.<sup>12</sup> The current gold standard for perforation identification is the use of a computerised tomographic scan.<sup>13,14</sup> While being widely available and helpful when radiation burden is a concern, ultrasonography makes pneumoperitoneum identification challenging, even for experienced sonographers.<sup>15,16</sup>

## 2. Materials and Methods

A prospective study of patients operated on for perforation peritonitis between March 1, 2021, and March 23, 2023 involved 104 patients with diagnosis of hollow viscus perforation with peritonitis in gastric and duodenal perforation were entered into this clinical study. Statistical comparisons were performed using the chi-square test and student's 't' test. Data was analyzed for Age wise distribution, Gender wise distribution.

**Inclusion criteria:** All cases of hollow viscus perforation with peritonitis in gastric and duodenal perforation from age 10 to 95 years.

**Exclusion criteria:** Any other perforation like illeal / appendicular.

**Table 1:** Age distribution

Age in Years	Frequency	Percentage
18-30 Yrs	25	20.8%
31-45 Yrs	24	20%
46-60 Yrs	23	19.1%
61-75 Yrs	20	16.6%
=/>76	12	10%
Total	104	100%

**Table 2:** Gender distribution

Gender	Number of Patients	Percentage
Male	81	78%
Female	23	22%
Total	104	100

## 3. Results


A total of 104 individuals had surgery for repair of gastroduodenal perforations. It was found that peptic ulcer perforation affects men more often than women since men outnumbered women by a ratio of 3.5 to 1. History of abdominal pain was present in every individual 104 in

**Table 3:** Modes of clinical presentation signs and symptoms

Fever	84	80.7 %
Nausea and Vomiting	100	96.1 %
Abdominal Pain	104	100 %
Rigors	54	51.9 %
Abdominal Distension	70	67.3 %
Hypotension	49	47.1%
Tachycardia	90	86.5 %
Guarding	99	95.1 %
Rigidity	100	96.1 %

(100%), vomiting was present in 100 in (96.1%) individuals. The other complaint were fever in 84 in (80.7%), rigors in 54 (51.9%), abdominal distension in 70 (67.3%), hypotension in 49 (47.1%), tachycardia in 90 (86.5%), guarding in 99 (91.1%), rigidity in 100 (96.1%) individuals. In addition, it was been discovered that younger age groups were more susceptible to peptic ulcer perforations. In comparison to gastric perforations, the mean age of duodenal perforations was 37.75, approximately 20 years less. Over 75 percent of duodenal perforations happened before the age of 50. We highlight that gastroduodenal ulcers share a similar pathophysiology, particularly with *H. pylori* infestation<sup>17</sup> which was more prevalent in younger patients from lower socioeconomic groups,<sup>17</sup> despite the fact that we do not have a good reason for this shifting epidemiological profile. Since up to three out of every five patients in this research had no prior dyspeptic history, it has been shown that perforation may be the earliest sign of peptic ulcer disease. While 43% of patients reported using some kind of antiulcer medicine in the six months before to perforation.

A dry fast is likely to result in a higher frequency of difficulties within a shorter time period from the start of fasting, as opposed to the partial hunger experienced during Ramadan. As 13.5% of our patients had ulcers bigger than 2 cm in diameter, our study has demonstrated that a repair using an omental patch or simple repair delivers satisfactory outcomes even for ulcers that are rather large. Five of the nine patients who had further surgeries following the treatment were determined to have an unbroken repair. At the time of the original operation, two patients developed fibrosis around the ulcer margin, and even after the ulcer margins were removed and an omental patch was pedicled, a leakage still occurred. In our series' overall mortality of 17.3% was within the range of 4-30% that is frequently cited in numerous series.<sup>18,19</sup> After operative management, two of our patients succumbed to pulmonary embolism resulted in 1.9% deaths, the rest from multiple organ failure, adult respiratory distress syndrome, and septicemia.



**Fig. 1:** X-ray chest showing continuous diaphragm sign and free air under diaphragm



**Fig. 3:** Intra op picture of Pre-pyloric perforation



**Fig. 2:** X-ray abdomen erect showing free air



**Fig. 4:** Intra oppicture of graham's omental patch repair

#### 4. Discussion

In this study, a total of 104 individuals had surgery to repair the gastroduodenal perforations. This results in an average of about 35 instances each year. When compared to other Eastern and Southern African series<sup>18</sup> and Enugu Nigeria, this number has a little higher prevalence. This may be an under representation, as numerous late cases may have passed away from the illness prior to receiving final surgery and are not included.

We discover that peptic ulcer perforation affects men more often than women since men outnumbered women by a ratio of 3.5 to 1. This result is in line with a number of previous African studies that demonstrate a male preponderance, ranging from a low of 1.3: 1 in Bugando, Tanzania<sup>19</sup> to a high ratios of 14.3:1 in Ido Ekiti, Nigeria<sup>18</sup> and 8.3:1 in Techiman, Ghana.<sup>16</sup> Contrary to popular belief, it is not a condition that only affects older females as shown in western television programmes.<sup>20,21</sup>

In addition to the aforementioned, it has been discovered that younger age groups were more susceptible to peptic ulcer perforations. In comparison to gastric perforations, the mean age of duodenal perforations was 37.75, approximately 20 years less. Over 75 percent of duodenal

perforations happened before the age of 50. The youngest patient in this series, it should be mentioned, had a stomach perforation while taking ibuprofen for a week to treat a severe low backache brought on by farm activity. Even in the past, this medication has been connected to peptic ulcers before in the paediatric age group.[32]

In contrast to other research from Nigeria that found no instances of gastric ulcer perforation,[27,30,34]<sup>22,23</sup> we now report that the ratio of gastric ulcers to duodenal perforations is almost 2:1. Several additional African studies demonstrate a majority of duodenal perforation, despite a prior research from a city hospital in Ghana having revealed a similar conclusion<sup>16</sup>. We highlight that gastroduodenal ulcers share a similar pathophysiology, particularly with *H. pylori* infestation,<sup>17</sup> which was more prevalent in younger patients from lower socioeconomic groups,<sup>24,25</sup> despite the fact that we do not have a good reason for this shifting epidemiological profile. Studies from Southern and Northern Nigeria show that urease culture tests for antral biopsies had a high incidence of 81.4% and over 90% with serological testing among individuals with dyspepsia. While *H. pylori* testing was not accessible in our facility for this investigation, all patients with perforated ulcers underwent *H. pylori* eradication medication. Additionally, overuse of nonsteroidal anti-inflammatory drugs, which we detected in as many as 37.5% of cases, is a significant etiologic factor, particularly in cases of stomach ulceration. Other risk variables were "dry" fasting and the use of herbal medicines that other workers have previously mentioned. Dry fasting is defined as not eating or drinking during the fast.

Since up to three out of every five patients in this research had no prior dyspeptic symptoms, it has been shown that perforation may be earliest sign of peptic ulcer disease. 43% of patients reported using any kind of antiulcer medicine in the six months before to perforation, as has been previously noted that the diagnosis of peptic ulcer illness is often made only after perforation.<sup>26</sup> This number is marginally higher than the 31% of perforations in individuals with established chronic peptic ulcer disease that were reported in Enugu, Nigeria.<sup>22</sup> When compared to an acute aggravation of a peptic ulcer, which may postpone final surgical management, this finding has the distinct benefit of raising the index of suspicion of perforation. However, with perforation, we see that discomfort remained constant throughout our series, occurring in 100% of instances and being followed by vomiting in 71% of them. This discovery is comparable to that made by Chalya et al.<sup>19</sup> who noted these two prominent presenting characteristics. Only 31% of our patients experience fever, making it a far less common symptom among our patients. The usage of analgesics with antipyretic effects may be to blame for this discovery.

Despite some patient were not willing or did not cooperate for computed tomographic scan at our centre

during the research period, this study demonstrated a high detection rate of pneumoperitoneum of 91%. This is more than other research' predictions<sup>11</sup> but comparable to what was discovered in Ghana<sup>16</sup> As lengthy intervals between perforation and radiologic examination increase radiographic identification of pneumoperitoneum, late presentation may be a factor.<sup>13</sup> Multidetector CT offers the specific benefit of directly demonstrating gastrointestinal discontinuity at the location<sup>26</sup> and aiding in diagnosis We also propose that simple radiographs, as other workers have stated,[31] are sufficient in our environment for emergency patients with rapid onset epigastric pain to help in choosing the best surgical choice for perforated peptic ulcers.

Naturally, the bulk of our patients belonged to lower socioeconomic classes in a rural town like ours. However, the clergy or pastors were another risk category identified by this study. Over a five-year span, four pastors who had perforated peptic ulcers had surgery. They were all male and had gastrointestinal perforations. Before they had perforation, they had been dry fasting for three to seven days.

Two other female patients, one with a stomach perforation and the other with a duodenal one, were also hospitalised with symptoms. Both admitted with symptoms while one who was fasting, suffered with stomach perforation. Peptic ulcers and their consequences are more common during the Ramadan fast, according to several studies in the past A dry fast is likely to result in a higher frequency of difficulties within a shorter time period from the start of fasting, as opposed to the partial hunger experienced during Ramadan.

As 13.5% of our patients had ulcers bigger than 2 cm in diameter, our study has demonstrated that a repair using an omental patch or simple repair delivers satisfactory outcomes even for ulcers that were rather large. Five of the nine patients who had further surgeries following the treatment were determined to have an unbroken repair. At the time of the original operation, two patients developed fibrosis around the ulcer margin, and even after the ulcer margins were removed and an omental patch was pedicled, a leakage still occurred. Our series' overall mortality of 17.3% is within the range of 4-30% that is frequently cited in numerous series.<sup>27</sup>

After reoperations, two of our patients succumbed. Pulmonary embolism resulted in 1.9% deaths, the rest from multiple organ failure, adult respiratory distress syndrome, and septicemia.

## 5. Conclusion

We state in conclusion that, the perforated peptic ulcer is a prevalent surgical issue in contemporary society. The majority of these perforations are gastroduodenal in nature, and for most patients, they constitute the initial indication of peptic ulcer disease. To determine the diagnosis in this case,

a simple chest radiograph in the classic case of sudden onset epigastric pain is enough.

## 6. Source of Funding

None.

## 7. Conflict of Interest

None.

## References

- Bertleff M, Lange JF. Perforated peptic ulcer disease: a review of history and treatment. *Dig Surg*. 2010;27(3):161–9.
- Lau JY, Sung J, Hill C, Henderson C, Howden CW, Metz DC, et al. Systematic review of the epidemiology of complicated peptic ulcer disease: incidence, recurrence, risk factors and mortality. *Digestion*. 2011;84(2):102–13.
- Williams N, Bullstrode C, O'connell P. Stomach and Duodenum in Bailey and Love's Short Practice of Surgery. 26th ed. and others, editor; 2013. p. 1539.
- Wang YR, Richter JE, Dempsey DT, Guzel H, Kahramanca S, Şeker D, et al. Peptic ulcer complications requiring surgery: what has changed in the last 50 years in Turkey. *Turk J Gastroenterol*. 1993;251(1):152–5.
- Milosavljevic T, Milosavljevi MK, Jovanovi I, krstic M. Complications of peptic ulcer disease. *Dig Dis*. 2011;29:491–3.
- Perforatedulcer; 2017. Available from: <http://en.m.wikipedia.org/wiki/perforatedulcer>.
- Hirschowitz BI, Simmons J, Mohnen J. Clinical outcome using lansoprazole in acid hypersecretors with and without Zollinger-Ellison syndrome: a 13-year prospective study. *Clin Gastroenterol Hepatol*. 2005;3(1):39–48.
- Fakhry SM, Watts DD, Luchette FA. Current diagnostic approaches lack sensitivity in the diagnosis of perforated blunt small bowel injury: analysis from 275,557 trauma admissions from the EAST multi-institutional HVI trial. *J Trauma Injury Infect Crit Care*. 2003;54(2):295–306.
- Saverio SD, Bassi M, Smerieri N. Diagnosis and treatment of perforated or bleeding peptic ulcers. *World J Emerg Surg*. 2000;6:9–45.
- Woodring JH, Heiser MJ. Detection of pneumoperitoneum on chest radiographs: comparison of upright lateral and posteroanterior projections. *Am J Roentgenol*. 1995;165(1):45–7.
- Baker SR, Maniatis V, Chryssikopoulos H, Roussakis A. Perforation of the alimentary tract: Evaluation with computed tomography. *Abdominal Imag*. 1996;21(5):373–9.
- Coppolino FF, Gatta G, Grezia GD. Gastrointestinal perforation: Ultrasonographic diagnosis. *Crit Ultrasound J*. 2013;5(1):S4. doi:10.1186/2036-7902-5-S1-S4.
- Wegdam HH, Hillah AA. Modified Open Omental Plugging of Peptic Ulcer Perforation in a Municipal Hospital in Ghana. *Postgrad Med J Ghana*. 2013;2(1):20–3.
- Ugochukwu AI, Amu OC, Nzegwu MA, Dilibe UC. Acute perforated peptic ulcer: on clinical experience in an urban tertiary hospital in south east Nigeria. *Internationalm J Surg*. 2013;11(3):223–7.
- Chalya PL, Mabula JB, Koy M. Clinical profile and outcome of surgical treatment of perforated peptic ulcers in northwestern Tanzania: a tertiary hospital experience. *World J Emerg Surg*. 2011;6:31. doi:10.1186/1749-7922-6-31.
- Shein M, Saadia R. Perforated peptic ulcer at the J.G Strijdom hospital: a retrospective study of 99 patients. *S Afr Med J*. 1986;70(5):21–3.
- Khan SB, Riaz N, Afza N. Perforated Peptic Ulcers: A review of 36 cases. *Professional Med J*. 2011;18:124–51.
- Oribhabor FO, Adebayo BO, Aladesanmi T. Akinola DO: Perforated duodenal Ulcer; Management in a resource poor, semi-urban Nigerian Hospital,” in rian Hospital. *Hospital J Surg*. 2013;19(1):13–5.
- Kang JY, Elders A, Majeed A, Maxwell JD, Bardhan KD. Recent trends in hospital admissions and mortality rates for peptic ulcer in Scotland 19822002. *Alimentary Pharmacology Therap*. 2006;24(1):65–79.
- Gona SK, Alassan MK, Marcellin KG. Postoperative Morbidity and Mortality of Perforated Peptic Ulcer: Retrospective Cohort Study of Risk Factors among Black Africans in Cote d'Ivoire. *Gastroent Res Pract*. 2016;p. 2640730. doi:10.1155/2016/2640730.
- Dakubo JCB, Naaader SB, Clegg-Lamprey JN. Gastroduodenal peptic ulceration. *East Afr med J*. 2009;86(3):100–9.
- Wang SY, Cheng CT, Liad CT, Wong HW, Cy YC. Surgical planning in Perforated peptic ulcer. *Medscape*;21:755–61.
- Williams S, O'connell PR, Mccaskie A. Bailey & Love's short practice of surgery. and others, editor. CRC press; 2022. p. 1088.
- Nuhu A, Kassama Y. Experience with acute perforated duodenal ulcer in a West African population. *Niger J Med*. 2008;17(4):403–6.
- Malfertheiner P, Chan FK, Mccoll KE. Peptic ulcer disease. *The Lancet*. 2009;374(9699):1449–61.
- Grassi R, Romano S, Pinto A, Romano L. Gastro-duodenal perforations: conventional plain film, US and CT findings in 166 consecutive patients. *Eur J Radiol*. 2004;50(1):30–6.
- Thorsen K, Glomsaker TB, Meer A, Soreide K, Soreide JA. Trends in diagnosis and surgical management of patients with perforated peptic ulcer. *J Gastrointest Surg*. 2011;15(8):1329–35.

## Author biography

**Pranjal Tanter**, Junior Resident

**Pratik Gond**, Senior Resident

**Shreya Pakkal**, Junior Resident

**Manideep Thippaarapu**, Junior Resident

**Kundankumar N Dandekar**, Professor

**Cite this article:** Tanter P, Gond P, Pakkal S, Thippaarapu M, Dandekar KN. Clinical study of 104 cases of hollow viscus perforation with peritonitis in gastric and duodenal perforation at rural tertiary centre. *IP J Surg Allied Sci* 2023;5(3):83-87.