



Original Research Article

Paper on role of cervical disc replacement surgery in cervical pvd patients in both elderly and young population and it's comparison with anterior cervical discectomy and fusion

Dhrumil Patel^{1*}, Dhruv Patel¹, Khanjan Ayar¹

¹Dept. of Orthopaedics, B.J. Medical College, Ahmedabad, Gujarat, India



ARTICLE INFO

Article history:

Received 20-11-2023

Accepted 21-12-2023

Available online 09-02-2024

Keywords:

Anterior cervical discectomy and fusion (ACDF); Arthrodesis

Arthroplasty

Artificial disc

Cervical disc arthroplasty (CDA)

Cervical spine

Cervical spondylosis

Disc prosthesis

Replacement

Fusion

Myelopathy

Radiculopathy

ABSTRACT

Introduction: For more than fifty years, anterior cervical discectomy and fusion (ACDF) has been a widely accepted operation for cervical spine degenerative diseases, including disc herniation and spondylosis, that are refractory to medical management. The high rates of success and patients' satisfaction with ACDF has made the operation a standard of care in common neurosurgical practice, however, the advantages of cervical disc arthroplasty (CDA) over fusion including maintaining normal neck motion and reducing degeneration of adjacent segments of the cervical spine

Aims and Objectives: To study the role of cervical disc replacement surgery in cervical pvd patients in both elderly and young population and it's comparison with anterior cervical discectomy and fusion with age variability taken into consideration.

Materials and Methods: The inclusion and exclusion criteria used, The operative procedure for both ACDF and CDA, The further follow up studies considering clinical as well as radiological outcome

Observation and Results: The most commonly indexed levels of CDA were C4–5 and C5–6, accounting for 85% of the entire series.

The clinical outcomes, including VAS of neck and arm pain, NDI, and JOA scores were all similarly satisfactory after the surgery, during the follow-up.

Conclusion: CDAs has been demonstrated to preserve segmental motion at long-term followup and to decrease the incidence of adjacent-level surgery, as compared to ACDF. Cervical global alignment and the segmental angle at rest improves after cervical arthroplasty

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

For more than fifty years, anterior cervical discectomy and fusion (ACDF) has been a widely accepted operation for cervical spine degenerative diseases, including disc herniation and spondylosis, that are refractory to medical management. The high rates of success and patients' satisfaction with ACDF has made the operation a standard of care in common neurosurgical practice, however, the

advantages of cervical disc arthroplasty (CDA) over fusion including maintaining normal neck motion and reducing degeneration of adjacent segments of the cervical spine.³ In the literature, CDA results are at least similar or even superior to clinical outcomes after anterior cervical arthrodesis at short- and medium-term follow-up. The implantation of CDA was reported to be a safe procedure, However, despite the low revision rates, favorable outcomes, feasibility, and ability to perform explanation of artificial disc prostheses, many surgeons still have negative perceptions of cervical arthroplasty. No study

* Corresponding author.

E-mail address: ahirkhanjan@yahoo.com (D. Patel).

that we are aware of has compared the adjacent-level reoperation rates for CDA versus the natural history of the disease. Furthermore, we believe that other significant factors limiting the more widespread use of CDA include the technical difficulty of the procedure compared to ACDF, complications such as spontaneous fusion and loosening, postoperative hematoma, heterotopic ossification (HO) and, in some countries, decreased reimbursement compared to ACDF. Moreover these reports have proven that CDA devices can maintain segmental mobility at the indexed level(s) and likely have the potential to reduce adjacent segment disease (ASD).¹

This paper is a review of the current literature, to determine the radiological and clinical outcomes of patients who underwent CDA for cervical degenerative disease and to compare with widely used ACDF procedure in cases of cervical myelopathy this in order to help inform preoperative decision-making and discussions with patients.

2. Materials and Methods

2.1. Inclusion & exclusion criteria

Consecutive adult patients (> 18 years of age) who underwent one- or two-level CDA with Prestige LP artificial discs (Medtronic, Memphis, TN) at the subaxial (C3–7) cervical spine in a single institute were included. The surgical indication for CDA was symptomatic cervical disc herniation and/or spondylosis causing radiculopathy, myelopathy, or both, that was refractory to medical treatment. All patients had failed at least 12 weeks of non-operative management, including physical and pain control therapy, but remained medically intractable prior to surgery.

2.2. Exclusion criteria were

Spinal trauma and fracture;^{1,2} evident segmental instability (i.e. more than 3.5mm translation or 20° angular motion);³ arthrodesis without mobility;⁴ severely incompetent facet joints;⁵ adjacent segment disease;⁶ ossification of posterior longitudinal ligament (OPLL);⁷ kyphotic deformity;⁸ infection; and⁹ long-term steroid use

2.3. Surgical technique

The standard Cloward approach for anterior cervical discectomy was executed in all patients. In addition to thorough discectomy, bilateral uncovertebral joints and bone spurs were removed extensively with drilling burrs or Kerrison's rongeurs to achieve generous decompression of the dura sac and nerve roots. Also, the posterior longitudinal ligaments were always resected to ensure adequate decompression. Upon placement of the Prestige LP artificial disc, meticulous endplate preparation, selection of a proper fitting size, and centering of the device were considered imperative to minimize the chances of

heterotopic ossification (HO) formation.² Furthermore, we used copious saline irrigation persistently to wash away the bone dust generated from osteophyte drilling in every case. All surgeries were done by three experienced neurosurgeons (JC Wu, WC Huang, and H Cheng) with consistent techniques detailed in our previous publications.^{3–8}




Figure 1:

2.4. Clinical and radiological follow up

Regular visits at the outpatient department were arranged at pre-operation, and post-operative 6-weeks, and at 3, 6, 12 and 24 months for all patients. Clinical follow-up parameters included visual analog scales (VAS), neck disability index (NDI), and modified Japanese Orthopaedic Association (JOA) scores, which were collected by experienced physician assistants during regular post-operative follow-ups. Routine X-ray images including antero-posterior, lateral, and flexion-extension films were taken at every regular visit at the clinic. Radiological criteria for adjacent segment degeneration (ASD) in X-ray films were the presence of disc space narrowing, osteophytes, or sclerosis of the endplates.⁹ Incidences of HO formation were assessed by not only the post-operative lateral radiographs but also by CT scans with three-dimensional reconstruction, and graded according to the McAfee's classification.¹⁰ Segmental range of motion (ROM) at the index level was Page 3 of 9 determined with standing lateral flexion/extension radiographs (Figure 1) at post-operative 24-months follow-up using the Cobb method.² Radiological measurements were completed by a board-certified neuroradiologist independently using the PACS system software, SmartIris (Taiwan Electronic Data Processing Co., Taiwan).

3. Observation and Results

A total of 171 patients underwent 1- or 2- level CDA with Prestige LP artificial discs, with an average age of 48.3±9.99 years at the time of operation. In order to




Figure 2:

investigate the age-related effects of CDA, the current study aimed to analyze patients at the two ends of the age distribution in the cohort. Therefore, the study included 71 patients who were aged 65 years or more and those less than 40 (to a minimum of 18 years), to minimize the bias from mid-aged CDA patients. The demographic data of these are demonstrated in Table 1. The patients were divided into two groups: the elderly group (≥ 65 years old) consisted of 24 patients with a mean age of 71.2 ± 4.79 years; the young group (≤ 40 years) composed of 47 patients with a mean age at 33.9 ± 4.45 year-old. The mean follow-up duration was 28.0 months, without differences between the two groups (averaged 22.6 versus 30.8 months, elderly vs. Young, respectively). The most commonly indexed levels of CDA were C4–5 and C5–6, accounting for 85% of the entire series. Approximately 50 min, consumed for the CDA surgery in the two age groups. The clinical outcomes, including VAS of neck and arm pain, NDI, and JOA scores

were all similarly satisfactory after the surgery, during the follow-up. All the patient-reported outcome parameters had significant improvement at the final follow-up, when compared to the pre-operative scores (Table 2).

4. Discussion

The current study focused on a comparison of patients at both ends of the age distribution (≥ 65 or ≤ 40 years) in a cohort of CDA patients. Patients who had been followed-up for more than 24 months were included for retrospective analysis of their clinical outcomes (i.e. VAS, NDI, and JOA scores), radiological parameters (i.e. pre-, post-operative ROM, and incidences of HO), and the complication profiles (e.g. dysphagia, dysphonia, and C5 palsy). The study demonstrated small but distinct discrepancies between the elderly and young in the changes of segmental mobility after CDA. More than 2 years after 1- or 2-level surgery for CDA, both groups of patients, regardless of their age differences (i.e. elderly versus young), demonstrated improvements in clinical outcomes when compared to that of pre-operation. Moreover, both groups had successfully preserved segmental mobility at the indexed levels with CDA. Comparing different prostheses with ACDF or between the various types of prostheses is reasonable. The clinical application of the Discover cervical artificial disc was first reported by Greiner–Perth et al.¹¹ in 2009. Discover prosthesis is both a metal-on-polymer type and a non-constrained type of prosthesis, which may have advantages in terms of the incidence of heterotopic ossification and ASD.

In our study, the postoperative neck disability index (NDI) and VAS scores of both the ACDF group and the TDR group were significantly improved from their preoperative scores, and there was no significant difference between the two groups at different time points (up to at least a two-year follow-up). Published randomised controlled clinical studies have shown the non-inferiority of short-term outcomes of TDR surgery compared to ACDF surgery.^{12–14} We similarly did not find a significant difference in overall complication rates due to adjacent segment degeneration between the two procedures at a mean of 5.4 years of follow-up across both groups.

Veeravagu et al.¹⁵ reviewed 28,777 cases undergoing ACDF and found that 9.13% of the single-level and 10.7% of the multilevel ACDF groups required a second operation within two years. They reported that the number of fusion levels was significantly correlated with the rate of reoperations. At 48 months, Davis et al.¹⁶ reported a cumulative incidence of 4% reoperation at the index level in the TDR group.

As more research is conducted on artificial disc replacements and surgical techniques continue to improve, it is likely that ADR will become an even more important part of spine care in the future. For patients suffering from

Table 1: Demographic data

Elderly group (265year -old)	Young group (S40 year -old)	P value
Case number n =24	n =47	
Age (years)71.2±4.79(65-80)	33.9±4.45(23-39)	<0.001*
Male :Female 13:11	31:16	
Follow -up (months)"226±22.29	308±21.52	0.145
Total levels 40	57	
one -level 8	37	
two -level 16	10	
Level distribution		0.253
	2	
C4-5 15	16	
C5-6 17	35	
C6-7 4	4	
Blood loss (ml)171.4±164.75	115.2±131.41	0.178
Operative time (min)"239.4±81.40	178.6±58.00	0.008*

"Values are presented as mean ±SD (standard deviation) *p <0.05;

Table 1 Demographic data

Elderly group (265year -old)	Young group (S40 year -old)	P value
Case number n =24	n =47	
Age (years)71.2±4.79(65-80)	33.9±4.45(23-39)	<0.001*
Male :Female 13:11	31:16	
Follow -up (months)"226±22.29	308±21.52	0.145
Total levels 40	57	
one -level 8	37	
two -level 16	10	
Level distribution		0.253
	2	
C4-5 15	16	
C5-6 17	35	
C6-7 4	4	
Blood loss (ml)171.4±164.75	115.2±131.41	0.178
Operative time (min)"239.4±81.40	178.6±58.00	0.008*

"Values are presented as mean ±SD (standard deviation) *p <0.05;

Table 2: Clinical outcomes

	Mean pr-op	Final follow-up(SD)	P value	Means pre-op Vas arm (SD)	Final follow-up (SD)	P value
Elderly	4.9(2.67)	2.0(1.70)	0.002*	4.5	1.9(2.07)	0.013*
Young group	3.1(2.68)	2.0(2.12)	0.020*	2.7(2.71)	1.0(1.75)	0.004*
	Mean pre-op NDI (SD)	Final follow-up (SD)	P value	Mean pre-op JOA (SD)	Final Follow-up(SD)	P value
Elderly group	19.2(12.44)	8.6(6.50)	0.012*	10.0(4.45)	13.14(2.79)	0.0009*
Young group	9.9(7.90)	6.0(4.34)	0.014*	13.3(1.67)	16.0 (1.19)	<0.001*

VAS Visual Analog scale for pain, NDI Neck Disability index, JOA Modified Japanese Orthopaedic Association score, SD standard deviation Pvalue: pair-ttest compared to pre-operative scores.*p<0.05

back pain or other spinal conditions, ADR offers a safe and effective alternative to traditional spinal surgery that can help them regain their mobility and quality of life.

This recent study highlights the benefits of cervical disc arthroplasty and reinforces the importance of considering all options when deciding on a treatment plan for cervical disc disease. Patients should discuss their individual needs and the risks and benefits of ADR with a board-certified physician to determine if it is right for them.

5. Conclusion

There are numerous benefits to CDAs. Cervical arthroplasty can mitigate pain and neck disability in appropriate patients with degenerative cervical disc disease causing radiculopathy or myelopathy. CDAs has been demonstrated to preserve segmental motion at long-term followup and to decrease the incidence of adjacent-level surgery, as compared to ACDF. Cervical global alignment and the segmental angle at rest improves after cervical arthroplasty.

On the other hand, there are also some negatives associated with CDA. It is not appropriate for anyone with osteoporosis, ossifying diseases, instability, collapsed disc, facet arthrosis, inflammatory diseases, spinal infections and retro vertebral disease. It is also technically demanding to implant a perfectly sized, perfectly placed prosthesis and there are numerous pitfalls that can result in poor outcomes. Despite the challenges, when performed technically well in appropriate patients, we believe that cervical arthroplasty is a safe and effective alternative to anterior cervical arthrodesis with several potential benefits.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

- Nunley PD, Coric D, Frank KA, Stone MB. Cervical Disc Arthroplasty: Current Evidence and Real-World Application. *Neurosurgery*. 2018;83(6):1087–106.
- Tu TH, Wu JC, Huang WC, Wu CL, Ko CC, Cheng H, et al. The effects of carpentry on heterotopic ossification and mobility in cervical arthroplasty: determination by computed tomography with a minimum 2-year follow-up: clinical article. *J Neurosurg Spine*. 2012;16(6):601–10.
- Tu TH, Huang WC, Tsai HW, Ko CC, Fay LY. Differences between 1 and 2-level cervical arthroplasty: more heterotopic ossification in 2-level disc replacement: Clinical article. *J Neurosurg Spine*. 2012;16(6):594–600.
- Fay LY, Huang WC, Tsai TY, Wu JC, Ko CC, Tu TH. Differences between arthroplasty and anterior cervical fusion in two-level cervical degenerative disc disease. *Eur Spine J*. 2014;23(3):627–61.
- Chang HK, Huang WC, Wu JC, Tu TH, Fay LY, Chang PY. Cervical arthroplasty for traumatic disc herniation: an age- and sex-matched comparison with anterior cervical discectomy and fusion. *BMC Musculoskelet Disord*. 2015;16:228–50.
- Chang HK, Huang WC, Wu JC, Chang PY, Tu TH, Fay LY. Should cervical disc arthroplasty be done on patients with increased intramedullary signal intensity on magnetic resonance imaging? *World Neurosurg*. 2016;89:489–96.
- Chang PY, Chang HK, Wu JC, Huang WC, Fay LY, Tu TH. Differences between C3-4 and other subaxial levels of cervical disc arthroplasty: more heterotopic ossification at the 5-year follow-up. *J Neurosurg Spine*. 2016;24(5):752–61.
- Chang HK, Chang CC, Tu TH, Wu JC, Huang WC, Fay LY. Can segmental mobility be increased by cervical arthroplasty? *Neurosurg Focus*. 2017;42(2):3–3.
- Hilibrand AS, Yoo JU, Carlson GD, Bohlman HH. The success of anterior cervical arthrodesis adjacent to a previous fusion. *Spine*. 1976;22(14):1574–83.
- Mcafee PC, Cunningham BW, Devine J, Williams E, Yu-Yahiro J. Classification of heterotopic ossification (HO) in artificial disk replacement. *J Spinal Disord Tech*. 2003;16(4):384–93.
- Perth RG, Allam Y, Silbermann J, Simank HG. First experience and preliminary clinical results with the cervical disc replacement DISCOVER. *Z Orthop Unfall*. 2009;147(5):582–9.
- Heller JG, Sasso RC, Papadopoulos SM. Comparison of BRYAN cervical disc arthroplasty with anterior cervical decompression and fusion: clinical and radiographic results of a randomized, controlled, clinical trial. *Spine*. 1976;34(2):101–7.
- Murrey D, Janssen M, Delamarter R, Goldstein J, Zigler J, Tay B, et al. Results of the prospective, randomized, controlled multicenter Food and Drug Administration investigational device exemption study of the ProDisc-C total disc replacement versus anterior discectomy and fusion for the treatment of 1-level symptomatic cervical disc disease. *Spine J*. 2009;9:275–86.
- Sasso RC, Anderson PA, Riew KD, Heller JG. Results of cervical arthroplasty compared with anterior discectomy and fusion: four-year clinical outcomes in a prospective, randomized controlled trial. *J Bone Joint Surg Am*. 2011;93(18):1684–92.
- Veeravagu A, Cole T, Jiang B, Ratliff JK. Revision rates and complication incidence in single- and multilevel anterior cervical discectomy and fusion procedures: an administrative database study. *Spine J*. 2014;14(7):1125–56.
- Hillips FM, Lee JY, Geisler FH. A prospective, randomized, controlled clinical investigation comparing PCM cervical disc arthroplasty with anterior cervical discectomy and fusion. 2-year results from the US FDA IDE clinical trial. *Spine*. 1976;38(15):E907–18.

Author biography

Dhruvil Patel, Senior Resident

Dhruv Patel, Senior Resident

Khanjan Ayar, Resident

Cite this article: Patel D, Patel D, Ayar K. Paper on role of cervical disc replacement surgery in cervical pvd patients in both elderly and young population and it's comparison with anterior cervical discectomy and fusion. *IP J Surg Allied Sci* 2023;5(4):114-118.